

Persistent basophilia may suggest an ‘accelerated phase’ in the evolution of CALR-positive primary myelofibrosis towards acute myeloid leukemia

Jerome Dobrowolski¹, Sergiu Pasca¹, Patric Teodorescu¹, Cristina Selicean², Ioana Rus², Mihnea Zdrengeha¹, Anca Bojan¹, Adrian Trifa¹, Bogdan Fetica², Ana Maria Rosu¹, Bobe Petrushev³, Ioana Berindan-Neagoe¹, Ciprian Tomuleasa^{1*}, Delia Dima²

¹Iuliu Hațieganu University of Medicine and Pharmacy, Romania, ²Oncology Institute Prof. Dr. Ion Chiricuta, Romania, ³Octavian Fodor Gastroenterology Institute, Romania

Submitted to Journal:
Frontiers in Oncology

Specialty Section:
Hematologic Malignancies

ISSN:
2234-943X

Article type:
Hypothesis and Theory Article

Received on:
07 Jun 2019

Accepted on:
21 Aug 2019

Provisional PDF published on:
21 Aug 2019

Frontiers website link:
www.frontiersin.org

Citation:
Dobrowolski J, Pasca S, Teodorescu P, Selicean C, Rus I, Zdrengeha M, Bojan A, Fetica B, Rosu A, Petrushev B, Berindan-neagoe I, Tomuleasa C and Dima D (2019) Persistent basophilia may suggest an ‘accelerated phase’ in the evolution of CALR-positive primary myelofibrosis towards acute myeloid leukemia. *Front. Oncol.* 9:872. doi:10.3389/fonc.2019.00872